

Chromosomal-level genome assembly of the painted sea urchin *Lytechinus pictus*, a genetically enabled model system for cell biology and embryonic development

Jacob F. Warner^{1*}, James W. Lord¹, Samantha A. Schreiter¹, Katherine T. Nesbit², Amro Hamdoun²,
Deirdre C. Lyons^{2*}

1. Department of Biology and Marine Biology, University of North Carolina Wilmington, 601 S. College Rd, Wilmington, NC 28403, USA

2. Scripps Institution of Oceanography, University of California San Diego, 9500 Gilman Dr, La Jolla, CA 92093-0202, USA

* Corresponding Authors:

Jacob Warner: warnerj@uncw.edu

Deirdre Lyons: d1lyons@ucsd.edu

Keywords: *Lytechinus pictus*, sea urchin, echinoderm, genome

Significance statement

The sea urchin *Lytechinus pictus* has long served as a model system for developmental, toxicological, and ecological studies. Here we present a chromosomal-level genome assembly as a resource for these communities and other researchers using this important model system.

© The Author(s) 2021. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Abstract

The painted urchin *Lytechinus pictus* is a sea urchin in the family Toxopneustidae and one of several sea urchin species that are routinely used as an experimental research organism. Recently, *L. pictus* has emerged as a tractable model system for establishing transgenic sea urchin lines due to its amenability to long term laboratory culture. We present the first published genome of *L. pictus*. This chromosomal-level assembly was generated using Illumina sequencing in conjunction with Oxford Nanopore Technologies long read sequencing and HiC chromatin conformation capture sequencing. The 998.9 Mb assembly exhibits high contiguity and has a scaffold length N50 of 46.0 Mb with 97% of the sequence assembled into 19 chromosomal-length scaffolds. These 19 scaffolds exhibit a high degree of synteny compared to the 19 chromosomes of a related species *Lytechinus variegatus*. *Ab initio* and transcript evidence gene modeling, combined with sequence homology, identified 28,631 gene models that capture 92% of BUSCO orthologs. This annotation strategy was validated by manual curation of gene models for the ABC transporter superfamily, which confirmed the completeness and accuracy of the annotations. Thus, this genome assembly, in conjunction with recent high contiguity assemblies of related species, positions *Lytechinus pictus* as an exceptional model system for comparative functional genomics and it will be a key resource for the developmental, toxicological, and ecological biology scientific communities.

Introduction

For over a century, biologists have used sea urchin eggs and embryos to make important contributions to the fields of development, cell biology, and gene regulation (Driesch, 1891; Evans et al., 1983; Foltz and Hamdoun, 2019; Hamdoun and Foltz, 2019; Oliveri et al., 2008). The attributes that make sea urchins excellent material for developmental and cell biology research include large quantities of sperm and oocytes, rapid development into pelagic larva with a relatively simple and well-defined set of cell types, and optically clear embryos and larvae that facilitate fluorescence microscopy and *in-vivo* imaging (Foltz and Hamdoun, 2019; Hamdoun and Foltz, 2019).

Unlike similar research carried out in other model systems, the communities using sea urchins have not converged on a singular species to work with, as researchers typically favor working with local species. For example, *Paracentrotus lividus* is used in Europe (Campos et al., 2020), *Heliocidaris tuberculata* is used in Australia (Morris et al., 2019), *Hemicentrotus pulcherrimus* is used in Asia (Liu et al., 2019), *Arbacia* spp. and *Lytechinus variegatus* are used on the east coast of North America (Hogan et al., 2020; Queiroz et al., 2021), and *Stronglyocentrotus purpuratus* and *Lytechinus pictus* are used on the west coast of North America (Nesbit and Hamdoun, 2020; Peter, 2019). The investment in many sea urchin species, instead of just one, has paid out in numerous studies comparing sea urchin (and echinoderm) development, gene regulatory networks, and genomes (Cary and Hinman, 2017; Erkenbrack and Thompson, 2019; Wang et al., 2020).

Although most sea urchin experimental material is collected from wild populations, a few sea urchin species are regularly bred in captivity including *Temnopleurus reevesii* (Yaguchi, 2019; Yaguchi et al., 2020) and *Lytechinus variegatus* (Heflin and Watts, 2016). Long before those studies, Ralph Hinegardner and colleagues (Cameron and Hinegardner, 1974; Coffaro and Hinegardner,

1977; Hinegardner, 1969) established inbred lines of *Lytechinus pictus*. This species is well-suited for rearing in the lab because it is small as an adult (2-4 cm in diameter), has a short generation time (~4-6 months), and is resistant to infection in captivity (Hinegardner, 1975). *L. pictus* is used for studying fertilization (Hussain et al., 2017), gastrulation (Aleksanyan et al., 2016), toxicology (Li et al., 2020), and response to ocean acidification (Smith et al., 2019), among other areas. Given the long history of research with *L. pictus*, our groups previously developed updated culturing methods, obtained developmental transcriptomes, and established a detailed staging system for this species (Nesbit et al., 2019; Nesbit and Hamdoun, 2020), with the ultimate goal of establishing inbred and germline transgenic lines.

A key impediment to functional and comparative genomics in *L. pictus* is the availability of a high contiguity genome. To remove this impediment, we have obtained and annotated a high-quality whole genome sequence for *L. pictus*. The *L. pictus* genome follows a recently published genome sequence in a sister species: *Lytechinus variegatus* (Davidson et al., 2020). The *L. pictus* genome is comparable to that of *L. variegatus* in size (998 Mb), chromosome number (likely 19), and chromosomal organization, as we show here. The completion of the *L. pictus* genome thus establishes the sea urchin as an excellent model system for comparative genomic studies as high contiguity assemblies are now available in several species representing a range of taxa (Davidson et al., 2020; Kinjo et al., 2018; Sodergren et al., 2006). Furthermore, the expanding genomics toolkit for *L. pictus* will maximize its usefulness for a wide range of research fields and support work on establishing stable transgenic sea urchin lines.

Results and Discussion

Lytechinus pictus genome assembly and synteny analysis

The *L. pictus* genome assembly, which we term *Lpic_2.0* (*Lpic_1.0* refers to an unpublished assembly), totals 998.85 Mb of sequences (0.10% gaps) assembled into 1307 scaffolds with half of the sequence (N50) contained in 8 scaffolds longer than 46 Mb (Table 1). 97% of the sequence is assembled into nineteen scaffolds longer than 10 MB, corresponding to putative chromosomes. This result is in agreement with a recent genome sequencing effort in *Lytechinus variegatus*, which also identified 19 chromosome-length scaffolds (Davidson et al., 2020). To identify homologous chromosomes between the two assemblies, we performed a nucleotide alignment between the 19 longest scaffolds in both species (Figure 1). *L. variegatus* and *L. pictus* diverged approximately 5-10 million years ago (Zigler and Lessios, 2004) and this is reflected in the genome alignment that shows a high overall level of synteny between *L. variegatus* and *L. pictus*, albeit with a large number of inversions and translocations, particularly in regard to chromosomes 1 and 5 (Figure 1).

Gene annotation

To predict protein coding genes, we used the MAKER pipeline (Cantarel et al., 2008). We used available protein models from *S. purpuratus* and *L. variegatus* in conjunction with a previously assembled transcriptome for homology and evidence-based gene modeling. We then used the resultant gene models to train Augustus and Snap for two additional rounds of *de novo* gene prediction. This resulted in 28,631 protein coding gene annotations, including 17,865 with 5'UTRs and 17,760 with 3'UTR sequences (Table 1, Supplementary Table 1). These protein coding gene models contained 92% of the Benchmarking Universal Single-Copy Orthologs (BUSCO) of metazoan genes (Simão et al., 2015). We also functionally annotated the gene models using protein alignments against the *S. purpuratus* protein, RefSeq protein, and Uniprot (Swissprot and Trembl) databases. 91% (26,179) of the gene models yielded a significant hit to the RefSeq database (blastp, e value $< 5e-5$) and 89% (25,483) yielded a significant hit to the *S. purpuratus* protein database.

Finally, we used the top hit from Uniprot to associate GO terms to our gene models. All of the annotations can be found in supplementary files 1 and 2.

Manual curation of ABC transporter annotations

To further assess the completeness and quality of the *L. pictus* genome, we manually examined the annotations of the ATP-Binding Cassette (ABC) transporter superfamily. This group of genes has a number of features which make it attractive for assembly assessment: The ABC transporters are a large multigene family (Dean and Annilo, 2005), they include large genes which can span several hundred kilobases (Luciani et al., 1994), and they are well characterized in a related echinoderm, *S. purpuratus*. The proteins are essential for ion regulation (Bryan et al., 2007), developmental signaling (Petrášek and Friml, 2009), cell migration (Kassmer et al., 2015), antigen presentation (Schumacher et al., 1994), and in humans, mutations in these genes are associated with disease (Supplementary Table 2) (Borst and Elferink, 2002; Dean, 2002; Gottesman and Ambudkar, 2001; Silverton et al., 2011; Tarling et al., 2013). There were 54 *L. pictus* ABC gene candidates in our assembly, spanning ABCs found in sea urchins (A-H) (Goldstone et al., 2006; Shipp and Hamdoun, 2012). These include putative orthologs of disease-relevant genes such as ABCA1, ABCB1, ABCC1, ABCC9, ABCD1 and ABCG5 (Supplementary Table 2). To assess the quality of individual annotations we focused on the ABCC subfamily (Supplementary Figure 2.). A feature of ABC families in the sea urchins is the relative expansion of the ABCC family as compared to humans (Gokirmak et al., 2016). Consistent with this, *L. pictus* have 20 putative C-subfamily candidates compared to the 12 ABCC members in human (Supplementary file 4). Of these C-subfamily members in urchins, the *S. purpuratus* ABCC1 contains 29 exons and spans approximately 45.5 kb. We identified a *L. pictus* ABCC1 candidate of 31 exons and 59.9 kb, encoding a 1,471 aa open reading frame. Similarly, the *S. purpuratus* ABCC4a gene is encoded by 27 exons spanning approximately 43.8 kb in length. A corresponding annotation for this isoform of ABCC4a in *L. pictus* is 24 exons and approximately

31.7 kb long, encoding a 1,334 aa open reading frame. Collectively these results support the outputs of the automated pipeline annotations, and the significant hits of *L. pictus* gene models against RefSeq and *S. purpuratus* protein databases, supporting the ‘completeness’ of our assembly and annotation.

This chromosomal-level genome assembly will be a valuable resource for the design of transgenic lines in this genetically enabled model system that is also well suited to laboratory culture through multiple generations (Hinegardner, 1969; Nesbit et al., 2019). Our annotation identifies homologues for the large majority of developmental regulatory genes described in the well-established sea urchin model *S. purpuratus*, thus enabling evolutionary comparisons between these two model systems. Furthermore, this genome assembly, in conjunction with the recent genome assembly of the sister species *Lytechinus variegatus* represent two of the most complete assemblies in any echinoderm model system. These chromosome-level assemblies will allow for robust synteny analyses to identify conserved cis-regulatory regions between the two species. Together, these two high-contiguity genomes firmly establish the sea urchin as an excellent model system for comparative genomics and represent a major advancement in the field for sea urchin developmental biology.

Materials and Methods

Tissue collection

Adult *L. pictus* were collected near the Scripps Institution of Oceanography, in La Jolla, CA and held in free-flowing sea water aquaria. Spawning was induced by injection of 0.5 M potassium chloride. From one of the males, several hundred microliters of sperm were “dry” collected using a pipette, centrifuged to remove excess seawater and flash frozen in liquid nitrogen and stored at -80°C. DNA extraction was carried out by Dovetail Genomics (Scott’s Valley, CA).

Dovetail Sequencing and assembly

A Chicago library was prepared as described previously (Putnam et al., 2016). Briefly, ~500ng of HMW gDNA (mean fragment length = 75kbp) was reconstituted into chromatin *in vitro* and fixed with formaldehyde. Fixed chromatin was digested with DpnII, the 5' overhangs filled in with biotinylated nucleotides, and free blunt ends ligated. After ligation, crosslinks were reversed and the DNA purified from protein. Purified DNA was treated to remove biotin that was not internal to ligated fragments. The DNA was then sheared to ~350 bp mean fragment size and sequencing libraries were generated using NEBNext Ultra enzymes and Illumina-compatible adapters. Biotin-containing fragments were isolated using streptavidin beads before PCR enrichment of each library. The libraries were sequenced on an Illumina HiSeq X platform to produce 387 million 2x150bp paired end reads, which provided 42.50 x physical coverage of the genome (1-100 kb pairs).

Two Dovetail HiC libraries were prepared in a similar manner as described previously (Lieberman-Aiden et al., 2009). Briefly, for each library, chromatin was fixed in place with formaldehyde in the nucleus and then extracted. Fixed chromatin was digested with DpnII, the 5' overhangs filled in with biotinylated nucleotides, and free blunt ends ligated. After ligation, crosslinks were reversed and the DNA purified from protein. Purified DNA was treated to remove biotin that was not internal to ligated fragments. The DNA was then sheared to ~350 bp mean fragment size and sequencing libraries were generated using NEBNext Ultra enzymes and Illumina-compatible adapters. Biotin-containing fragments were isolated using streptavidin beads before PCR enrichment of each library. The libraries were sequenced on an Illumina HiSeq X platform. The number and length of read pairs produced for each library was: 247 million, 2x150bp for library 1 and 170 million, 2x150bp for library 2. Together, these Dovetail HiC library reads provided 2,771.85 x physical coverage of the genome (10-10,000 kb pairs, supplementary figure 1).

Five runs of ONT MinION sequencing generated 15 Gb of data representing 19X coverage. The ONT data was assembled with the wtdbg2 assembler, followed by consensus polishing with ONT reads with racon and Illumina polishing with pilon. The input *de novo* assembly, ONT reads, Chicago library reads, and Dovetail HiC library reads were used as input data for HiRise, a software pipeline designed specifically for using proximity ligation data to scaffold genome assemblies (Putnam et al., 2016). An iterative analysis was conducted. First, Shotgun and Chicago library sequences were aligned to the draft input assembly using a modified SNAP read mapper (<http://snap.cs.berkeley.edu>). The separations of Chicago read pairs mapped within draft scaffolds were analyzed by HiRise to produce a likelihood model for genomic distance between read pairs, and the model was used to identify and break putative misjoins, to score prospective joins, and make joins above a threshold. After aligning and scaffolding Chicago data, Dovetail HiC library sequences were aligned and scaffolded following the same method. After scaffolding, shotgun sequences were used to close gaps between contigs.

Synteny analysis

The 19 largest scaffolds were repeat masked using RepeatFinder and aligned to the 19 chromosomes identified in the *L. variegatus* genome using D-Genies (Cabanettes and Klopp, 2018) with minimap as the aligner to identify homologous chromosomes. The *L. pictus* scaffolds were renamed for the homology to *L. variegatus* (e.g. “chr1”), with the remaining chromosomes enumerated and named with the prefix “unplaced_scaffold”.

Annotation

Genome annotation was performed using the MAKER pipeline (Cantarel et al., 2008). A repeat library was generated with RepeatModeler and repeats were identified and masked prior to

annotation (Smit and Hubley 2010). MAKER was run using protein models from *S. purpuratus* (available on echinobase: http://legacy.echinobase.org/Echinobase/download/downloadfile.php?file=SPU_peptide.fasta.zip ; last accessed 3/5/20) and *L. variegatus* (available on echinobase: <http://legacy.echinobase.org/Echinobase/download/downloadfile.php?file=Lvar2.2-protein.fa.zip>; last accessed 3/5/20) along with a previously assembled transcriptome from *L. pictus* (Nesbit et al., 2019) for a first round of homology-based and RNA-evidenced gene prediction. The resultant gene models were used for two subsequent rounds of training and *ab initio* gene prediction using SNAP and Augustus as part of the MAKER pipeline. The gene models were annotated using the notation LPI_XXXXXX and analyzed for ‘completeness’ with BUSCO version 4 using the metazoan gene set (Simão et al., 2015). Functional annotation was performed using BLASTp to compare our gene models to the *S. purpuratus* protein database, RefSeq database, and Uniprot (Swissprot and TrEMBL) databases (hits defined as having an e value < 5e-5). The top hit to the Uniprot database was used to source putative GO terms via programmatic retrieval from uniprot.org.

ABC Transporter annotation

To identify putative ABC transporters, known sequences from human, mouse, and *S. purpuratus* were used as input for BLAST searches of the *L. pictus* genome and transcriptome. Redundant sequences and low-quality hits were eliminated. To help confirm the identity of the *L. pictus* ABC candidates we analyzed the amino acid sequences of the deduced protein. This included manual notation of transmembrane domains, and the Walker A/B/C domains, which are characteristic of ABCs, using Geneious (v11.1.5). Candidate *L. pictus* ABC transporters that had strong support from our initial BLAST search and contained the predicted structural motifs were further validated with a reciprocal BLAST approach. Each of our candidate sequences served as the query in a BLAST search against annotated genes and proteins in the NCBI, SwisProt, and Echinobase databases. We

excluded hypothetical and uncharacterized proteins. Hits were deemed true ABC transporter orthologs if supported with a combination of high (>85%) sequence identity to known ABC transporters, structural motif analysis, and reciprocal BLAST search. To annotate beyond subfamily resolution, we focused on the ABCC candidates from *L. pictus* and performed phylogenetic analysis to verify individual genes. Sequences were aligned in Geneious (v11.1.5) using the ClustalW program with default settings (gap open = 10.0, gap extension = 0.1, word size = 1, window length = 5, topdiag = 5, pairgap = 3). The resultant alignment was run through ProtTest (v3.4.2) to predict the best fit model for tree construction. A maximum likelihood tree (RaxML-HPC2 on XSEDE) with 1000 bootstraps was then produced using the LG+G model, with yeast selected as an outgroup, through the CIPRES Science Gateway (v3.3) (Miller et al., 2011). Remaining parameters were used on default settings. The resulting tree was exported and formatted using FigTree (V1.4.4).

Data Availability

The sequence assembly and annotations can be found at the National Center for Biotechnology Information (NCBI) under Bioproject PRJNA647794. Additionally, annotation files including a GFF3 and functional annotation file can be found in supplementary files 1 and 2.

Acknowledgements

This work was supported by NIH ES 027921 and 030318, and NSF 1840844 to AH, by a Scripps Institution of Oceanography SEED Grant to DCL and AH, and by a U.C. San Diego Academic Senate Grant to DCL.

References

Aleksanyan, H., Liang, J., Metzzenberg, S., Oppenheimer, S.B., 2016. Terminal alpha-d-mannosides are critical during sea urchin gastrulation. *Zygote* 24, 775-782.

Borst, P., Elferink, R.O., 2002. Mammalian ABC transporters in health and disease. *Annual Review of Biochemistry* 71, 537-592.

Bryan, J., et al. 2007. ABCC8 and ABCC9: ABC transporters that regulate K⁺ channels. *Pflugers Archiv-European Journal of Physiology* 453, 703-718.

Cabanettes, F., Klopp, C., 2018. D-GENIES: dot plot large genomes in an interactive, efficient and simple way. *PeerJ* 6, e4958-e4958.

Cameron, R.A., Hinegardner, R.T., 1974. Initiation of metamorphosis in laboratory cultured sea urchins. *Biol Bull* 146, 335-342.

Campos, S., Troncoso, J., Paredes, E., 2020. Major challenges in cryopreservation of sea urchin eggs. *Cryobiology*. S0011-2240(20)30652-0.

Cantarel, B.L., et al. 2008. MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome research* 18, 188-196.

Cary, G.A., Hinman, V.F., 2017. Echinoderm development and evolution in the post-genomic era. *Developmental Biology* 427, 203-211.

Coffaro, K.A., Hinegardner, R.T., 1977. Immune response in the sea urchin *Lytechinus pictus*. *Science* 197, 1389-1390.

Davidson, P.L., et al. 2020. Chromosomal-Level Genome Assembly of the Sea Urchin *Lytechinus variegatus* Substantially Improves Functional Genomic Analyses. *Genome Biology and Evolution* 12, 1080-1086.

Dean, M., 2002. The Human ATP-Binding Cassette (ABC) Transporter Superfamily. National Center for Biotechnology Information (US), Bethesda, Maryland.

Dean, M., Annilo, T., 2005. Evolution of the ATP-binding cassette (ABC) transporter superfamily in vertebrates. *Annual Review of Genomics and Human Genetics* 6, 123-142.

Driesch, H., 1891. "Entwicklungsmechanische Studien: I. Der Werthe der beiden ersten Furchungszellen in der Echinogdermenentwicklung. Experimentelle Erzeugung von Theil- und Doppelbildungen. II. Über die Beziehungen des Lichte zu der ersten Etappe der thierischen Formbildung." *Zeitschrift für wissenschaftliche Zoologie* 53: 160–84. Translated as "The Potency of the First Two Cleavage Cells in Echinoderm Development. Experimental Production of Partial and Double Formations." In B. H. Willier and J. M. Oppenheimer (eds.), *Foundations of Experimental Embryology*. Hafner Press, New York, 38–50.

Erkenbrack, E.M., Thompson, J.R., 2019. Cell type phylogenetics informs the evolutionary origin of echinoderm larval skeletogenic cell identity. *Commun Biol* 2, 160.

Evans, T., Rosenthal, E.T., Youngblom, J., Distel, D., Hunt, T., 1983. Cyclin: A protein specified by maternal mRNA in sea urchin eggs that is destroyed at each cleavage division. *Cell* 33, 389-396.

Foltz, K.R., Hamdoun, A., 2019. *Methods in Cell Biology: Echinoderms, Part A, First Edition* ed. Academic Press, Cambridge, MA, United States.

Gokirmak, T., et al. 2016. Functional diversification of sea urchin ABCC1 (MRP1) by alternative splicing. *American Journal of Physiology-Cell Physiology* 310, C911-C920.

Goldstone, J.V., et al. 2006. The chemical defense: Environmental sensing and response genes in the *Strongylocentrotus purpuratus* genome. *Developmental Biology* 300, 366-384.

Gottesman, M.M., Ambudkar, S.V., 2001. Overview: ABC transporters and human disease. *Journal of Bioenergetics and Biomembranes* 33, 453-458.

Hamdoun, A., Foltz, K.R., 2019. *Methods in Cell Biology: Echinoderms Part B, First ed.* Academic Press, Cambridge, MA, United States.

Heflin, L.E., Watts, S.A., 2016. Feeding time, frequency and ration affect growth and energy allocation in young adults of the sea urchin *Lytechinus variegatus*. *Aquaculture Nutrition* 22, 1055-1064.

Hinegardner, R.T., 1969. Growth and development of the laboratory cultured sea urchin. *Biol Bull* 137, 465-475.

Hinegardner, R.T., 1975. Morphology and genetics of sea urchin development. *American Zoologist* 15, 679-689.

Hogan, J.D., et al. 2020. The developmental transcriptome for *Lytechinus variegatus* exhibits temporally punctuated gene expression changes. *Developmental Biology* 460, 139-154.

Hussain, Y.H., Sadilek, M., Salad, S., Zimmer, R.K., Riffell, J.A., 2017. Individual female differences in chemoattractant production change the scale of sea urchin gamete interactions. *Dev Biol* 422, 186-197.

Kassmer, S.H., Rodriguez, D., Langenbacher, A.D., Bui, C., De Tomaso, A.W., 2015. Migration of germline progenitor cells is directed by sphingosine-1-phosphate signalling in a basal chordate. *Nature Communications* 6.

Kinjo, S., Kiyomoto, M., Yamamoto, T., Ikeo, K., Yaguchi, S., 2018. HpBase: A genome database of a sea urchin, *Hemicentrotus pulcherrimus*. *Dev Growth Differ* 60, 174-182.

Li, A., Espinoza, J., Hamdoun, A., 2020. Inhibitory effects of neurotoxin β -N-methylamino-L-alanine on fertilization and early development of the sea urchin *Lytechinus pictus*. *Aquat Toxicol* 221, 105425.

Lieberman-Aiden, E., et al. 2009. Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* 326, 289-293.

Liu, D.M., Awazu, A., Sakuma, T., Yamamoto, T., Sakamoto, N., 2019. Establishment of knockout adult sea urchins by using a CRISPR-Cas9 system. *Development Growth & Differentiation* 61, 378-388.

Luciani, M.F., Denizot, F., Savary, S., Mattei, M.G., Chimini, G., 1994. Cloning of 2 novel ABC transporters mapping on human chromosome 9. *Genomics* 21, 150-159.

Miller, M.A., Pfeiffer, W., Schwartz, T., 2011. The CIPRES Science Gateway: A Community Resource for Phylogenetic Analyses. *Proceedings of the 2011 TeraGrid Conference: Extreme Digital Discovery*, 1-8. DOI 10.1145/2016741.2016785

Morris, V.B., Kable, E., Koop, D., Cisternas, P., Byrne, M., 2019. Early development of the feeding larva of the sea urchin *Heliocidaris tuberculata*: role of the small micromeres. *Development Genes and Evolution* 229, 1-12.

Nesbit, K.T., et al. 2019. The painted sea urchin, *Lytechinus pictus*, as a genetically-enabled developmental model. *Methods Cell Biol* 150, 105-123.

Nesbit, K.T., Hamdoun, A., 2020. Embryo, larval, and juvenile staging of *Lytechinus pictus* from fertilization through sexual maturation. *Dev Dyn* 249, 1334-1346.

Oliveri, P., Tu, Q., Davidson, E.H., 2008. Global regulatory logic for specification of an embryonic cell lineage. *Proceedings of the National Academy of Sciences of the United States of America* 105, 5955-5962.

Peter, I.S., 2019. Chapter 2 - Methods for the experimental and computational analysis of gene regulatory networks in sea urchins, in: Hamdoun, A., Foltz, K.R. (Eds.), *Methods in Cell Biology*. Academic Press, pp. 89-113.

Petrášek, J., Friml, J., 2009. Auxin transport routes in plant development. *Development* 136, 2675-2688.

Putnam, N.H., et al. 2016. Chromosome-scale shotgun assembly using an in vitro method for long-range linkage. *Genome research* 26, 342-350.

Queiroz, V., Muxel, S.M., Inguglia, L., Chiaramonte, M., Custódio, M.R., 2021. Comparative study of coelomocytes from *Arbacia lixula* and *Lythechinus variegatus*: Cell characterization and in vivo evidence of the physiological function of vibratile cells. *Fish & Shellfish Immunology* 110, 1-9.

Schumacher, T.N.M., et al. 1994. Peptide length and sequence specificity of the mouse TAP1/TAP2 translocator. *Journal of Experimental Medicine* 179, 533-540.

Shipp, L.E., Hamdoun, A., 2012. ATP-binding cassette (ABC) transporter expression and localization in sea urchin development. *Developmental Dynamics* 241, 1111-1124.

Silverton, L., Dean, M., Moitra, K., 2011. Variation and evolution of the ABC transporter genes ABCB1, ABCC1, ABCG2, ABCG5 and ABCG8: implication for pharmacogenetics and disease. *Drug Metabolism and Drug Interactions* 26, 169-179.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., Zdobnov, E.M., 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31, 3210-3212.

Smit AFA, Hubley R, Green P. RepeatMasker Open-4.0. 2013–2015. Available from: <http://www.repeatmasker.org>. Accessed February 6, 2020.

Smit AFA, Hubley R. RepeatModeler Open-1.0. 2008–2015. Available from: <http://www.repeatmasker.org>. Accessed February 6, 2020.

Smith, K.E., et al. 2019. Sea urchin reproductive performance in a changing ocean: poor males improve while good males worsen in response to ocean acidification. *Proc Biol Sci* 286, 20190785.

Sodergren, E., et al. 2006. The genome of the sea urchin *Strongylocentrotus purpuratus*. *Science* 314, 941-952.

Tarling, E.J., Vallim, T.Q.D., Edwards, P.A., 2013. Role of ABC transporters in lipid transport and human disease. *Trends in Endocrinology and Metabolism* 24, 342-350.

Wang, L., et al. 2020. Genetic basis for divergence in developmental gene expression in two closely related sea urchins. *Nat Ecol Evol* 4, 831-840.

Yaguchi, S., 2019. *Temnopleurus* as an emerging echinoderm model. *Methods Cell Biol* 150, 71-79.

Yaguchi, S., et al. 2020. Establishment of homozygous knock-out sea urchins. *Curr Biol* 30, R427-r429.

Zigler, K.S., Lessios, H.A., 2004. Speciation on the coasts of the new world: phylogeography and the evolution of *bindin* in the sea urchin genus *Lytechinus*. *Evolution* 58, 1225-1241.




Fig 1: Dot plot alignment of *L. variegatus* chromosomal sequences to the 19 largest *L. pictus* scaffolds (putative chromosomes). Dot plot was generated using D-Genies (Cabanettes and Klopp, 2018) with minimap as the alignment engine. The top 100,000 matches are displayed.

Assembly

Size	998.85 Mb
Scaffolds	1307
N50 Scaffold Length	46.003Mb
Longest Scaffold	91.80Mb
Scaffolds > 10MB	19
Sequence in 19 largest scaffolds	97%
N50 Contig Length	219.60Kb
N (%)	0.10
GC (%)	36

BUSCO

Complete	251
Duplicated	13
Fragmented	14
Missing	25

Annotations

# Protein Coding Genes	28631
# Gene Models with <i>Sp</i> hit	25483
# Genes Models with RefSeq hit	26179

Table 1: *Lytechinus pictus* genome assembly statistics




Fig 1: Dot plot alignment of *L. variegatus* chromosomal sequences to the 19 largest *L. pictus* scaffolds (putative chromosomes). Dot plot was generated using D-Genies (Cabanettes and Klopp, 2018) with minimap as the alignment engine. The top 100,000 matches are displayed.